

	Page
<b>Table 9.9.</b> Modified Bessel Functions—Orders 3–9 ( $0 \leq x \leq 20$ ) . . .	423
$e^{-x}I_n(x), e^xK_n(x), n=3(1)9$	
$x=0(.2)10(.5)20, 5S$	
<b>Table 9.10.</b> Modified Bessel Functions—Orders 10, 11, 20 and 21	
$(0 \leq x \leq 20)$ . . . . . 425	
$x^{-10}I_{10}(x), x^{-11}I_{11}(x), x^{10}K_{10}(x)$	
$x=0(.2)10, 8S$ or $9S$	
$e^{-x}I_{10}(x), e^{-x}I_{11}(x), e^xK_{10}(x)$	
$x=10(.2)20, 10D, 10D, 7D$	
$x^{-20}I_{20}(x), x^{-21}I_{21}(x), x^{20}K_{20}(x)$	
$x=0(.2)20, 5S$ to $7S$	
Modified Bessel Functions—Auxiliary Table for Large	
Arguments ( $20 \leq x \leq \infty$ ). . . . . 427	
$\ln\{x^{\frac{1}{2}}e^{-x}I_{10}(x)\}, \ln\{x^{\frac{1}{2}}e^{-x}I_{11}(x)\}, \ln\{\pi^{-1}x^{\frac{1}{2}}e^xK_{10}(x)\}$	
$\ln\{x^{\frac{1}{2}}e^{-x}I_{20}(x)\}, \ln\{x^{\frac{1}{2}}e^{-x}I_{21}(x)\}, \ln\{\pi^{-1}x^{\frac{1}{2}}e^xK_{20}(x)\}$	
$x^{-1}=.05(-.001)0, 8D, 6D$	
<b>Table 9.11.</b> Modified Bessel Functions—Various Orders ( $0 \leq n \leq 100$ ) .	428
$I_n(x), K_n(x), n=0(1)20(10)50, 100$	
$x=1, 2, 5, 10, 50, 100, 9S$ or $10S$	
<b>Table 9.12.</b> Kelvin Functions—Orders 0 and 1 ( $0 \leq x \leq 5$ ) . . . . .	430
ber $x$ , bei $x$ , ber <sub>1</sub> $x$ , bei <sub>1</sub> $x$	
ker $x$ , kei $x$ , ker <sub>1</sub> $x$ , kei <sub>1</sub> $x$	
$x=0(.1)5, 10D, 9D$	
Kelvin Functions—Auxiliary Table for Small Arguments	
$(0 \leq x \leq 1)$ . . . . . 430	
ker $x + \text{ber } x \ln x, \text{kei } x + \text{bei } x \ln x$	
$x(\text{ker}_1 x + \text{ber}_1 x \ln x), x(\text{kei}_1 x + \text{bei}_1 x \ln x)$	
$x=0(.1)1, 9D$	
Kelvin Functions—Modulus and Phase ( $0 \leq x \leq 7$ ) . . . . . 432	
$M_0(x), \theta_0(x), M_1(x), \theta_1(x)$	
$N_0(x), \phi_0(x), N_1(x), \phi_1(x)$	
$x=0(.2)7, 6D$	
Kelvin Functions—Modulus and Phase for Large Argu-	
ments ( $6.6 \leq x \leq \infty$ ). . . . . 432	
$x^{\frac{1}{2}}e^{-x/\sqrt{2}}M_0(x), \theta_0(x) - (x/\sqrt{2}), x^{\frac{1}{2}}e^{-x/\sqrt{2}}M_1(x), \theta_1(x) - (x/\sqrt{2})$	
$x^{\frac{1}{2}}e^{x/\sqrt{2}}N_0(x), \phi_0(x) + (x/\sqrt{2}), x^{\frac{1}{2}}e^{x/\sqrt{2}}N_1(x), \phi_1(x) + (x/\sqrt{2})$	
$x^{-1}=.15(-.01)0, 5D$	

---

The author acknowledges the assistance of Alfred E. Beam, Ruth E. Capuano, Lois K. Cherwinski, Elizabeth F. Godefroy, David S. Liepman, Mary Orr, Bertha H. Walter, and Ruth Zucker of the National Bureau of Standards, and N. F. Bird, C. W. Clenshaw, and Joan M. Felton of the National Physical Laboratory in the preparation and checking of the tables and graphs.